

FIAN TD-3

March 1994

AVERAGE MULTIPLICITIES IN GLUON AND QUARK JETS IN HIGHER-ORDER PERTURBATIVE QCD

I.M. Dremin¹ and V.A. Nechitailo²

*P.N.Lebedev Physical Institute,
Leninsky prospect,53, Moscow 117924, Russia.*

Abstract

The ratio of average multiplicities in gluon and quark jets is shown to become noticeably smaller in higher-order QCD compared to its lowest order value what improves agreement with experiment. QCD anomalous dimension has been calculated. It has been used to get energy dependence of mean multiplicities.

¹ e-mail address : dremin@td.fian.free.net

²e-mail address : nechit@td.fian.free.net

The ratio of average multiplicities in gluon ($\langle n_G \rangle$) and quark ($\langle n_F \rangle$) jets (let us call it $r \equiv \langle n_G \rangle / \langle n_F \rangle$) was of much debate during last years. Its value $N_c/C_F = 9/4$ predicted in the lowest order approximation of QCD is too large to fit experimental data. The simplest correction [1, 2] has reduced this value to 2.05. Recently it has been shown [3] that the exact solution of fixed coupling QCD equations gives rise to ever smaller value $r = 1.84 \pm 0.02$. The above values refer to the ratio of the number of partons in jets. In Monte Carlo computation using definite hadronization schemes they are recalculated to hadronic ratios observed in experiment which are about 40 percents smaller. Using one of such models (namely, Herwig), it has been shown in [3] that the above partonic value of r agrees quite well with experimental [4] hadronic value $1.27 \pm 0.04 \pm 0.06$.

We show here that quite similar values are obtained for the partonic ratio in the running coupling QCD if higher order terms are taken into account. Therefore, one concludes that this ratio is rather insensitive whether the coupling constant is fixed or running but requires proper treatment of energy conservation.

We have calculated also the QCD anomalous dimension in the same approximation and have found the energy dependence of average multiplicity. They are more sensitive to the running property of the coupling constant.

Let us remind briefly main notations. The generating functions of gluon $G_G(y, z)$ and quark $G_F(y, z)$ jets are

$$G_G(y, z) = \sum_{n=0}^{\infty} (1+z)^n P_n^G(y), \quad G_F(y, z) = \sum_{n=0}^{\infty} (1+z)^n P_n^F(y), \quad (1)$$

where $P_n^G(y)$ and $P_n^F(y)$ are the parton multiplicity distributions in gluon and quark jets, correspondingly, $y = \ln(Q/Q_0)$, Q is the jet virtuality, $Q_0 = \text{const}$. The normalized factorial moments F_q and Φ_q are related to generating functions as:

$$G_G = \sum_{q=0}^{\infty} \frac{z^q}{q!} \langle n_G \rangle^q F_q, \quad G_F = \sum_{q=0}^{\infty} \frac{z^q}{q!} \langle n_F \rangle^q \Phi_q, \quad (2)$$

The generating functions satisfy the following integro-differential equations [5]

$$G'_G(y) = \int_0^1 dx K_G^G(x) \gamma_0^2 [G_G(y + \ln x) G_G(y + \ln(1-x)) - G_G(y)] \\ + n_f \int_0^1 dx K_G^F(x) \gamma_0^2 [G_F(y + \ln x) G_F(y + \ln(1-x)) - G_G(y)], \quad (3)$$

$$G'_F(y) = \int_0^1 dx K_F^G(x) \gamma_0^2 [G_G(y + \ln x) G_F(y + \ln(1-x)) - G_F(y)], \quad (4)$$

where $G'(y) = dG/dy$, $\gamma_0^2 = 2N_c\alpha_S/\pi$, $\alpha_S = 2\pi/(11 - \frac{2}{3}n_f)y$ is the running coupling constant, n_f is the number of active flavours of quarks.

The kernels of equations are given by QCD as :

$$K_G^G(x) = \frac{1}{x} - (1-x)[2 - x(1-x)], \quad (5)$$

$$K_G^F(x) = \frac{1}{4N_c} [x^2 + (1-x)^2], \quad (6)$$

$$K_F^G(x) = \frac{C_F}{N_c} \left(\frac{1}{x} - 1 + \frac{x}{2} \right), \quad (7)$$

where $N_c = 3$ is the number of colours and $C_F = (N_c^2 - 1)/2N_c = 4/3$.

In solving eqs.(3),(4) we follow the method developed in [6, 7] and expand the generating functions in Taylor series at large y . We leave aside all terms with second and higher derivatives here because their contribution to average multiplicities is of higher order in coupling constant as can be easily checked. In that case, eqs.(3),(4) become

$$(\ln G_G(y))' = \int_0^y dy' \gamma_0^2(y') [G_G(y') - 1] \\ - h_1 \gamma_0^2 [G_G(y) - 1] + h_2 \gamma_0^2 G'_G(y) \\ + \frac{n_f}{12N_c} \gamma_0^2 \left\{ 2 \left[\frac{G_F^2(y)}{G_G(y)} - 1 \right] - h_4 \frac{G_F^2(y)}{G_G(y)} (\ln G_F(y))' \right\}, \quad (8)$$

$$(\ln G_F(y))' = \frac{C_F}{N_c} \left\{ \int_0^y dy' \gamma_0^2(y') [G_G(y') - 1] \right. \\ \left. + \gamma_0^2 \left[-\frac{3}{4} (G_G(y) - 1) + \frac{7}{8} G'_G(y) - h_3 (\ln G_F(y))' G_G(y) \right] \right\}, \quad (9)$$

where $h_1 = 11/24$, $h_2 = (67 - 6\pi^2)/36$, $h_3 = (4\pi^2 - 15)/24$, $h_4 = 13/3$.

To calculate average multiplicities, one should keep only the terms with $q = 0$ and 1 in (2). Normalization condition is $F_0 = F_1 = \Phi_0 = \Phi_1 = 1$. After differentiation, one gets from (8),(9)

$$\begin{aligned} \langle n_G \rangle'' = \gamma_0^2 \Big\{ & (1 + 4Bh_1\gamma_0^2)\langle n_G \rangle - 2h_1\langle n_G \rangle' + h_2\langle n_G \rangle'' \\ & + \frac{n_f}{12N_c} [2(2\langle n_F \rangle' - \langle n_G \rangle') - h_4\langle n_F \rangle'' \\ & - 4B\gamma_0^2(2\langle n_F \rangle - \langle n_G \rangle)] \Big\}, \end{aligned} \quad (10)$$

$$\langle n_F \rangle'' = \frac{C_F}{N_c} \gamma_0^2 \left\{ \left(1 + \frac{3}{2}B\gamma_0^2\right)\langle n_G \rangle - \frac{3}{4}\langle n_G \rangle' + \frac{7}{8}\langle n_G \rangle'' - h_3\langle n_F \rangle'' \right\}, \quad (11)$$

where $\gamma_0' = -B\gamma_0^3$, $B \equiv (\frac{11}{3}N_c - \frac{2}{3}n_f)/8N_c$ has been substituted. The QCD anomalous dimension $\gamma(y)$ is defined as

$$\langle n_G \rangle = \exp\left(\int^y \gamma(y')dy'\right). \quad (12)$$

We represent γ and r expanding them in γ_0 as

$$\gamma = \gamma_0(1 - a_1\gamma_0 - a_2\gamma_0^2) + O(\gamma_0^4), \quad (13)$$

$$r = \frac{N_c}{C_F}(1 - r_1\gamma_0 - r_2\gamma_0^2) + O(\gamma_0^3). \quad (14)$$

Since $\gamma_0' \sim \gamma_0^3$, we get $r' = 0$ up to terms $O(\gamma_0^3)$. However, such terms are taken into account in the formulae below. Up to correction terms of order $O(\gamma_0^2)$ one gets

$$\frac{\langle n_G \rangle'}{\langle n_G \rangle} = \gamma_0 [1 - a_1\gamma_0 - a_2\gamma_0^2], \quad (15)$$

$$\frac{\langle n_G \rangle''}{\langle n_G \rangle} = \gamma_0^2 [1 - (2a_1 + B)\gamma_0 - (2a_2 - a_1^2 - 2a_1B)\gamma_0^2], \quad (16)$$

$$\frac{\langle n_F \rangle'}{\langle n_G \rangle} = \frac{C_F}{N_c} \gamma_0 [1 + (r_1 - a_1)\gamma_0 + (r_2 + r_1^2 - a_2 - a_1r_1 - Br_1)\gamma_0^2], \quad (17)$$

$$\frac{\langle n_F \rangle''}{\langle n_G \rangle} = \frac{C_F}{N_c} \gamma_0^2 \left[1 + (r_1 - B - 2a_1)\gamma_0 + (r_2 + r_1^2 + a_1^2 - 3Br_1 - 2a_1r_1 - 2a_2 + 2Ba_1)\gamma_0^2 \right]. \quad (18)$$

We insert (15)-(18) into (10),(11) and keep again the terms of order $O(\gamma_0^2)$. Therefrom we obtain after simple algebraic manipulations

$$a_1 = h_1 + \frac{n_f}{12N_c^3} - \frac{B}{2}, \quad (19)$$

$$r_1 = 2 \left(h_1 + \frac{n_f}{12N_c^3} \right) - \frac{3}{4}, \quad (20)$$

$$a_2 = -\frac{1}{2} \left\{ h_2 - a_1^2 - 2a_1 \left(B - \left[h_1 + \frac{n_f}{12N_c^3} (1 + 4C_F N_c) \right] \right) - \frac{n_f}{12N_c} \frac{C_F}{N_c} [h_4 + 3 - 4B] + 4B \left[h_1 + \frac{n_f}{12N_c^3} \right] \right\}, \quad (21)$$

$$r_2 = \frac{r_1}{6} \left(\frac{25}{8} - \frac{3n_f}{4N_c} - \frac{C_F n_f}{N_c N_c} \right) + \frac{7}{8} - h_2 - \frac{C_F}{N_c} h_3 + \frac{n_f}{12N_c} \frac{C_F}{N_c} h_4. \quad (22)$$

Thus, we have obtained the QCD anomalous dimension and the ratio of average multiplicities in gluon and quark jets as functions of the QCD coupling constant γ_0 and of the number of active flavours n_f in higher-order perturbative QCD. The higher order corrections in r are negative and its value becomes smaller. The above dependences are very mild that agrees with conclusions of [3]. Besides, here we can insert the running coupling γ_0 in (13) and show dependence of γ on Q directly (see Fig.1). To do that, we use (as in [3]) the parameter $Q_0 = 0.65\Lambda_{\overline{\text{MS}}}$ with $\Lambda_{\overline{\text{MS}}} = 175\text{MeV}$ for $n_f = 5$ in proportion 63 : 100 : 130 for $n_f = 5 : 4 : 3$, correspondingly. Integration of these values of γ gives rise according to (12) to the following energy dependence of the average multiplicity

$$\langle n_G \rangle = y^{\frac{a_1}{2B}} \exp \left(\sqrt{\frac{2}{B}} y + \frac{a_2}{\sqrt{2B^3 y}} \right) \quad (23)$$

or in terms of γ_0 one obtains

$$\langle n_G \rangle = y^{\frac{a_1}{2B}} \exp[2\gamma_0(y)(1 + a_2\gamma_0^2(y))y] . \quad (24)$$

First term in the exponent corresponds to DLA-approximation with running coupling constant. There is an important difference between the fixed and running coupling constant impact on the energy behaviour of average multiplicities. Even though the fixed coupling case provides faster asymptotical increase of average multiplicity, in the energy range of interest up to $y \sim 3y_{Z^0}$ the mean multiplicity for fixed coupling increases slower than for running coupling. (Here $y_{Z^0} = 6.67$ is the scale at which the value of the fixed coupling has been calculated). It is shown in Fig.2.

The physical origin of the effect is easy to understand because rather small value of the fixed coupling has been used throughout the whole evolution of the cascade while in real situation it should become larger and larger at later stages what produces more particles in case of running coupling at lower energies. However, the multiplicity is enlarged if more flavours become active. It compensates somehow the lower values of the coupling constant at high virtualities what should be taken into account when comparison with experiment is done.

The preexponential term in expressions (23),(24) is also well known (see, e.g., [5, 9, 10]) and stems from MLLA-approximation³. The next-to-next-to leading correction to the exponent calculated by us diminishes it and the energy increase of the average multiplicity becomes somewhat slower (see Fig.3).

In Table 1 we show the values of the ratio r for various n_f at $Q = 91$ GeV (Z^0 mass). They are very close to those obtained for the fixed coupling QCD [3] and are about 20 percents smaller than the prediction of the lowest order approximation. We would like to point out that the γ_0^2 -correction to r calculated above is several times larger than its value found in [10] where the complicated diagram calculations have been done. The result of [10] is reproduced by the first term in (22). Other terms correspond to the contribution of second derivatives in (10),(11). For convenience, the numerical values of a_i, r_i are shown in the Table also.

Let us remind that one should combine these results with models for the conversion of partons into hadrons (hadronization) to get an access

³ Calculations in the framework of this approximation see, for example, in refs.[11, 12].

to experimental data. That is why, in ref.[3], the results of Monte Carlo calculation according to Herwig model [8] have been used. They show that at Z^0 -mass the ratio r at parton level exceeds its value at hadron level by about 40 percents. Being taken into account, this fact leads to agreement between theory and experiment [8]. Since our results are very similar to those of [3] we do not foresee any contradiction between them and experimental data.

To conclude, we have obtained the QCD anomalous dimension γ and the ratio of average multiplicities in gluon and quark jets in higher-order perturbative QCD with running coupling constant. The value of the ratio agrees rather well with the one previously obtained in fixed coupling QCD [3] by exact solution of the equations. Thus, it shows that the running property of the coupling constant is not as important for the ratio as proper treatment of higher-order terms including conservation laws but it is more noticeable in the energy dependence of mean multiplicity. Once again, it demonstrates that the values of r are much smaller than their lowest order counterparts and agree with experiment if the hadronization is properly taken into account. Nevertheless, one may still worry that the hadronization stage gives rise to the most important factor reducing parton ratios to their hadron values accessible in experiment. However, to get it in QCD one must solve the confinement problem that is out of our scope nowadays.

Acknowledgements

We are grateful to R.Hwa for valuable comments which help correct some formulae and improve the content.

This work was supported in part by Russian Fund for Fundamental Studies (grant 93-02-3815), by NATO grant CRG-930025 and by International Science Foundation grant M5V000.

References

- [1] A.H. Mueller, Nucl. Phys. **B241** (1984) 141.
- [2] E.D. Malaza and B.R. Webber, Phys. Lett. **B149** (1984) 501.
- [3] I.M. Dremin and R.C. Hwa, Preprint OITS 525 (1993), Phys. Lett. **B324** (1994) 477.
- [4] OPAL collaboration, P.D. Acton et al. Zs. Phys. **C58** (1993) 387.
- [5] Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller and S.I. Troyan, Basics of Perturbative QCD (Editions Frontieres, 1991).
- [6] I.M. Dremin, Phys. Lett. **B313** (1993) 209.
- [7] I.M. Dremin and V.A. Nechitailo, JETP Lett. **58** (1993) 945.
- [8] J.W. Gary, Preprint UCRHEP-T116 (1993).
- [9] F. Cuypers and K. Tesima, Zs. Phys. **C52** (1991) 69.
- [10] J.B. Gaffney and A.H. Mueller, Nucl. Phys. **B250** (1985) 109.
- [11] S. Catani , Yu.L. Dokshitzer , F. Fiorani and B.R. Webber , Nucl. Phys. **B377** (1992) 445.
- [12] Yu.L. Dokshitzer and M. Olsson, Nucl. Phys. **B396** (1993) 137.

Figure Captions

Fig. 1. The dependence of the anomalous dimension γ on Q . Solid lines are for γ and dashed lines are for γ_0 . The numbers of active flavours are indicated near lines in all figures.

Fig. 2. The y -dependence of the average multiplicity. Solid lines are for running coupling and dashed lines are for fixed coupling. The arrow marks the y_{Z^0} location.

Fig. 3. The y -dependence of the average multiplicity for the running coupling. Solid lines are for next-to-next-to leading approximation considered in the present paper and dashed lines are for MLLA.

Table 1:

n_f	r	r_1	r_2	γ_0	a_1	a_2
3	1.84	0.185	0.426	0.473	0.280	- 0.379
4	1.80	0.191	0.468	0.481	0.297	- 0.339
5	1.77	0.198	0.510	0.484	0.314	- 0.301

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ex/9406002v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ex/9406002v1>